



Mark scheme (Unused)

January 2022

Pearson Edexcel International Advanced Level
In Physics (WPH15/01)

Paper 5: Thermodynamics, Radiation, Oscillations
and Cosmology

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Autumn 2020

Publications Code xxxxxxxx*

All the material in this publication is copyright

© Pearson Education Ltd 2020

General Marking Guidance


- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Question Number	Answer	Mark
1	<p>B is the correct answer</p> <p>A is not correct, as binding energy is not related to temperature</p> <p>C is not correct, as a high collision rate is determined by the density</p> <p>D is not correct, as a very high temperature doesn't mean a high density</p>	(1)
2	<p>C is the correct answer, as $g = \frac{GM}{r^2}$ and $M = \rho V$</p>	(1)
3	<p>B is the correct answer</p> <p>A is not correct, as the electronic charge has been used to convert the mass</p> <p>C is not correct, as the conversion of mass to kg is incorrect</p> <p>D is not correct, as the conversion of mass to kg is incorrect</p>	(1)
4	<p>D is the correct answer,</p> <p>A is incorrect, as the amplitude decreases over time for a damped oscillation</p> <p>B is incorrect, as the amplitude stays constant over time for a free oscillation</p> <p>C is incorrect, as the amplitude may stay constant or decrease over time for a natural oscillation</p>	(1)
5	<p>D is the correct answer, as the count rate halves for each thickness of 1.5 cm</p>	(1)
6	<p>C is the correct answer, as $v_{\max} = 2\pi fA$</p>	(1)
7	<p>A is the correct answer, as $I = \frac{L}{4\pi d^2}$ gives d, so I and L must be known</p>	(1)
8	<p>B is the correct answer, as it is incorrect to say frequency decreases over time</p>	(1)
9	<p>B is the correct answer</p> <p>A is incorrect, as X is a diagram for a very old star cluster (white dwarf stars present)</p> <p>C and D are incorrect, as Z is a diagram for a medium age star cluster (red giant, but no white dwarf stars present)</p>	(1)
10	<p>C is the correct answer</p> <p>A is incorrect, as there are for more nucleons in a nucleus of ^{238}U than in ^{120}Sn</p> <p>B is incorrect, as a nucleus of ^{120}Sn has a higher B.E./nucleon than ^{16}O</p> <p>D is incorrect, as we cannot deduce this statement from the graph.</p>	(1)

Question Number	Answer	Mark
11	<p>Use of $V = \frac{4}{3}\pi r^3$ (1)</p> <p>Use of $pV = NkT$ (1)</p> <p>Conversion of temperature to kelvin (1)</p> <p>$N = 6.76 \times 10^{23}$ (1)</p> <p><u>Example of calculation</u></p> <p>$V = \frac{4}{3}\pi (0.185 \text{ m})^3 = 2.65 \times 10^{-2} \text{ m}^3$</p> <p>$N = \frac{pV}{kT} = \frac{1.04 \times 10^5 \text{ Pa} \times 2.65 \times 10^{-2} \text{ m}^3}{1.38 \times 10^{-23} \text{ J K}^{-1} \times (273 + 22.5) \text{ K}} = 6.76 \times 10^{23}$</p>	4
	Total for question 11	4

Question Number	Answer	Mark
12	<p>Use of $\frac{v}{c} = \frac{\Delta\lambda}{\lambda}$ (1)</p> <p>Use of $v = H_0 d$ (1)</p> <p>$d = 1.4 \times 10^{24} \text{ m}$ (1)</p> <p><u>Example of calculation</u></p> $v = \frac{\Delta\lambda}{\lambda} c = \frac{(438.6-434.1) \times 10^{-9} \text{ m}}{434.1 \times 10^{-9} \text{ m}} \times 3.00 \times 10^8 \text{ m s}^{-1} = 3.11 \times 10^6 \text{ m s}^{-1}$ $d = \frac{3.11 \times 10^6 \text{ m s}^{-1}}{2.3 \times 10^{-18} \text{ s}^{-1}} = 1.35 \times 10^{24} \text{ m}$	3
	Total for question 12	3

Question Number	Answer	Mark
13	<p>MAX 4</p> <p>The student is correct to say that the rate of decay decreases over time (1)</p> <p>However, the uranium doesn't become more stable, the number of unstable uranium nuclei decreases. (1)</p> <p>The student should have said that radiation is emitted from the nucleus [accept atom] (1)</p> <p>The student was wrong to say that the particles emitted are radioactive (1)</p> <p>Because the emitted particles do not decay (1)</p> <p>In a time equal to the half-life the number of unstable nuclei (and not the mass) decreases by 50%. (1)</p> <p>Because the product nuclei are nearly as massive as the unstable nuclei (1)</p>	4
	Total for question 13	4

Question Number	Answer	Mark
14(a)	<p>The star is viewed from two positions at 6 month intervals Or the star is viewed from opposite ends of the diameter of the Earth's orbit about the Sun (1)</p> <p>The change in angular position of the star against backdrop of distant/fixed stars is measured (1)</p> <p>Trigonometry is used to calculate the distance to the star [Do not accept Pythagoras] (1)</p> <p>The diameter/radius of the Earth's orbit about the Sun must be known (1)</p> <p>Full marks may be obtained from a suitably annotated diagram</p>  <p>[Accept the symmetrical diagram seen in many text books]</p>	4
14(b)	<p>Use of $s = ut$ (1)</p> <p>$s = 9.7 \times 10^{16} \text{ (m)}$ (1)</p> <p><u>Example of calculation</u></p> <p>$s = 3.00 \times 10^8 \text{ m s}^{-1} \times 10.3 \times (365 \times 86400) \text{ s} = 9.74 \times 10^{16} \text{ m}$</p>	2
Total for question 14		6

Question Number	Answer	Mark
15(a)	<p>Use of $E_k = \frac{1}{2}mv^2$ (1)</p> <p>Use of $E = mc\Delta\theta$ (1)</p> <p>Use of efficiency = $\frac{\text{useful energy output}}{\text{total energy input}}$ (1)</p> <p>Required energy = 1.9×10^4 J Or temperature rise = 29 K Or number of hits = 850 (1)</p> <p>Conclusion based on calculated values of energy transfer (1) Or Conclusion based on calculated value of temperature rise Or Conclusion based on calculated value of number of hits [conclusion must include a comparison of appropriate data]</p> <p><u>Example of calculation</u></p> <p>$E_k = \frac{1}{2} \times 1.1 \text{ kg} \times (7.5 \text{ m s}^{-1})^2 = 30.9 \text{ J}$</p> <p>$E = 1000 \times 30.9 \text{ J} \times 0.72 = 2.23 \times 10^4 \text{ J}$</p> <p>$E = (1.1 + 0.45) \text{ kg} \times 490 \text{ J kg}^{-1} \text{ K}^{-1} \times 25 \text{ K} = 1.90 \times 10^4 \text{ J}$</p> <p>[If mass of hammer neglected, $E = 0.45 \text{ kg} \times 490 \text{ J kg}^{-1} \text{ K}^{-1} \times 25 \text{ K} = 5.51 \times 10^3 \text{ J}$</p>	5
15(b)	<p>No thermal energy is transferred (from the steel plate) to surroundings Or hammer comes to rest after hitting the steel plate (1)</p> <p>[Allow no energy is used to deform the steel]</p>	1
	Total for question 15	6

Question Number	Answer	Mark
16(a)	<p>Use of $\omega = \frac{2\pi}{T}$ (1)</p> <p>Use of $v = r\omega$ (1)</p> <p>$v = 1.02 \times 10^3 \text{ m s}^{-1}$ (1)</p> <p><u>Example of calculation</u></p> <p>$\omega = \frac{2\pi}{27.3 \times 86400 \text{ s}} = 2.66 \times 10^{-6} \text{ rad s}^{-1}$</p> <p>$v = 3.84 \times 10^8 \text{ m} \times 2.66 \times 10^{-6} \text{ rad s}^{-1} = 1023 \text{ m s}^{-1}$</p>	3
16(b)(i)	<p>$\Delta E_{\text{grav}} = mg\Delta h$ is appropriate for situations in which g is approximately constant (1)</p> <p>As the distance moved is only a small fraction of the distance to the Earth, the value of g hardly changes (1)</p>	2
16(b)(ii)	<p>Use of $g = \frac{GM}{r^2}$ (1)</p> <p>Use of $\Delta E_{\text{grav}} = mg\Delta h$ (1)</p> <p>$\Delta E_{\text{grav}} = 7.6 \times 10^{19} \text{ J}$ (1)</p> <p>OR</p> <p>Use of $V_{\text{grav}} = -\frac{GM}{r}$ (1)</p> <p>Recognises that $\Delta E_{\text{grav}} = m \times \Delta V_{\text{grav}}$ (1)</p> <p>$\Delta E_{\text{grav}} = 7.6 \times 10^{19} \text{ J}$ (1)</p> <p><u>Example of calculation</u></p> <p>$g = \frac{6.67 \times 10^{-11} \text{ N m}^2 \text{ kg}^{-2} \times 6.02 \times 10^{24} \text{ kg}}{(3.84 \times 10^8 \text{ m})^2} = 2.72 \times 10^{-3} \text{ N kg}^{-1}$</p> <p>$\Delta E_{\text{grav}} = 7.35 \times 10^{22} \text{ kg} \times 2.72 \times 10^{-3} \text{ N kg}^{-1} \times 0.38 \text{ m} = 7.61 \times 10^{19} \text{ J}$</p>	3
	Total for question 16	8

Question Number	Answer	Mark																																								
*17	<p>This question assesses a student’s ability to show a coherent and logically structured answer with linkages and fully-sustained reasoning.</p> <p>Marks are awarded for indicative content and for how the answer is structured and shows lines of reasoning.</p> <p>The following table shows how the marks should be awarded for structure and lines of reasoning.</p> <table><tr><td></td><td>Number of marks awarded for structure of answer and sustained line of reasoning</td></tr><tr><td>Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning demonstrated throughout</td><td>2</td></tr><tr><td>Answer is partially structured with some linkages and lines of reasoning</td><td>1</td></tr><tr><td>Answer has no linkages between points and is unstructured</td><td>0</td></tr></table> <p>Total marks awarded is the sum of marks for indicative content and the marks for structure and lines of reasoning</p> <table><tr><th>IC points</th><th>IC mark</th><th>Max linkage mark</th><th>Max final mark</th></tr><tr><td>6</td><td>4</td><td>2</td><td>6</td></tr><tr><td>5</td><td>3</td><td>2</td><td>5</td></tr><tr><td>4</td><td>3</td><td>1</td><td>4</td></tr><tr><td>3</td><td>2</td><td>1</td><td>3</td></tr><tr><td>2</td><td>2</td><td>0</td><td>2</td></tr><tr><td>1</td><td>1</td><td>0</td><td>1</td></tr><tr><td>0</td><td>0</td><td>0</td><td>0</td></tr></table> <p>Indicative content</p> <p>IC1 Gravitational forces act on objects with mass</p> <p>IC2 But electric fields exert forces on objects with charge</p> <p>IC3 Electric forces can be attractive or repulsive, but gravitational forces are always attractive Or electric potential can be positive or negative, but gravitational potential is always negative.</p> <p>IC4 Both fields have an infinite range Or both fields are an example of “action at a distance”</p> <p>IC5 Both the gravitational field around a point mass and the electric field around a point charge obey an inverse square law</p> <p>IC6 The interaction between unit charges is larger than the interaction between unit masses (at a given separation) Or the electric force between point charges is larger than the gravitational force between unit masses (at a given separation).</p> <p>[for 2 linkage marks there must be at least 1 similarity and 1 difference]</p> <p>Total for question 17</p>		Number of marks awarded for structure of answer and sustained line of reasoning	Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning demonstrated throughout	2	Answer is partially structured with some linkages and lines of reasoning	1	Answer has no linkages between points and is unstructured	0	IC points	IC mark	Max linkage mark	Max final mark	6	4	2	6	5	3	2	5	4	3	1	4	3	2	1	3	2	2	0	2	1	1	0	1	0	0	0	0	6
	Number of marks awarded for structure of answer and sustained line of reasoning																																									
Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning demonstrated throughout	2																																									
Answer is partially structured with some linkages and lines of reasoning	1																																									
Answer has no linkages between points and is unstructured	0																																									
IC points	IC mark	Max linkage mark	Max final mark																																							
6	4	2	6																																							
5	3	2	5																																							
4	3	1	4																																							
3	2	1	3																																							
2	2	0	2																																							
1	1	0	1																																							
0	0	0	0																																							
		6																																								

Question Number	Answer	Mark
18(a)(i)	<p>Use of $P = \frac{\Delta E}{\Delta t}$ (1)</p> <p>Use of $\Delta E = c^2 \Delta m$ (1)</p> <p>$\frac{\Delta m}{\Delta t} = 5.04 \times 10^9 \text{ (kg s}^{-1}\text{)}$ (1)</p> <p><u>Example of calculation</u></p> <p>$\frac{\Delta m}{\Delta t} = \frac{4.54 \times 10^{26} \text{ W}}{(3.00 \times 10^8 \text{ m})^2} = 5.04 \times 10^9 \text{ kg s}^{-1}$</p>	3
18(a)(ii)	<p>Use of 0.08 % (1)</p> <p>Use of $\frac{\Delta m}{\Delta t}$ from (a) (1)</p> <p>$t = 9.9 \times 10^9 \text{ years}$ (ecf from (i)) (1)</p> <p><u>Example of calculation</u></p> <p>$\Delta m = \frac{0.08}{100} \times 1.97 \times 10^{30} \text{ kg} = 1.576 \times 10^{27}$</p> <p>$t = \frac{1.576 \times 10^{27} \text{ kg}}{5.04 \times 10^9 \text{ kg s}^{-1}} = 3.13 \times 10^{17} \text{ s} = 9.90 \times 10^9 \text{ years}$</p>	3
18(b)	<p>(Gamma Pavonis is more massive so) there is a greater temperature (and pressure) in the core (1)</p> <p>Rate of fusion is (much) higher than in delta Pavonis (1)</p> <p>Hence the time spent on main sequence is less and the suggestion is incorrect (1)</p> <p>MP3 dependent on MP2</p>	3
	Total for question 18	9

Question Number	Answer	Mark
19(a)(i)	Use of $\Delta F = k\Delta x$ with $F = mg$ (1) $k = 213 \text{ (N m}^{-1}\text{)}$ (1) <u>Example of calculation</u> $k = \frac{mg}{\Delta x} = \frac{65.0 \text{ kg} \times 9.81 \text{ N kg}^{-1}}{(48.0 - 45.0) \text{ m}} = 212.6 \text{ N m}^{-1}$	2
19(a)(ii)	(For simple harmonic motion the) acceleration is: <ul style="list-style-type: none"> (directly) proportional to displacement from equilibrium position (1) acceleration is in the opposite direction to displacement Or (always) acting towards the equilibrium position (1) OR (For simple harmonic motion the resultant) force is: <ul style="list-style-type: none"> (directly) proportional to displacement from equilibrium position (1) force is in the opposite direction to displacement Or (always) acting towards the equilibrium position (1) 	2
19(a)(iii)	Use of $T = 2\pi\sqrt{\frac{m}{k}}$ with $f = \frac{1}{T}$ (1) $f = 0.27 \text{ (Hz)}$ (1) <u>Example of calculation</u> $f = \frac{1}{2\pi}\sqrt{\frac{k}{m}} = \frac{1}{2\pi}\sqrt{\frac{210 \text{ N m}^{-1}}{75 \text{ kg}}} = 0.266 \text{ Hz}$	2
19(a)(iv)	Use of $\omega = 2\pi f$ (1) Use of $a = -\omega^2 x$ (1) $a = 3.4 \text{ m s}^{-2}$ (ecf from (iii)) (1) <u>Example of calculation</u> (1) $\omega = 2\pi \times 0.266 \text{ s}^{-1} = 1.67 \text{ rad s}^{-1}$ $a = (1.67 \text{ rad s}^{-1})^2 \times 1.2 \text{ m} = 3.35 \text{ m s}^{-2}$	3
19(b)	Work is done against air resistance (1) Or air resistance causes damping (1) So energy is transferred to the surroundings (1) Amplitude decreases to zero	3
Total for question 19		12

Question Number	Answer	Mark															
20(a)	<p>Top line correct (1) Bottom line correct (1)</p> ${}^{210}_{82}\text{Pb} \rightarrow {}^{210}_{83}\text{Bi} + {}^0_{-1}\beta^- + {}^0_0\bar{\nu}_e$	2															
20(b)	<p>Use of $\lambda = \frac{\ln 2}{t_{1/2}}$ (1) Use of $\frac{\Delta N}{\Delta t} = (-)\lambda N$ (1) Use of $A = A_0 e^{-\lambda t}$ (1) Activity is 25 Bq after 1.87 years, so claim is false. (1)</p> <p><u>Example of calculation:</u></p> $\lambda = \frac{\ln 2}{t_{1/2}} = \frac{0.693}{372 \times 86\,400 \text{ s}} = 2.16 \times 10^{-8} \text{ s}^{-1}$ $\frac{\Delta N}{\Delta t} = \lambda N = 2.16 \times 10^{-8} \text{ s}^{-1} \times 4.15 \times 10^9 = 89.6 \text{ Bq}$ $25 = 89.6 \times e^{-2.16 \times 10^{-8} t}$ $-2.16 \times 10^{-8} \text{ s}^{-1} \times t = \ln\left(\frac{25 \text{ Bq}}{89.6 \text{ Bq}}\right)$ $t = \frac{-1.28}{-2.16 \times 10^{-8} \text{ s}^{-1}} = 5.91 \times 10^7 \text{ s} = 1.87 \text{ year}$	4															
20(c)	<p>One pair of readings taken from graph and Rx^2 calculated (1) 2 more pairs of readings taken from graph and Rx^2 calculated (1) Check if Rx^2 is constant and conclusion consistent with calculations (1)</p> <p><u>Example of calculation</u></p> <table border="1"> <thead> <tr> <th>R / s^{-1}</th><th>x / cm</th><th>$Rx^2 / \text{cm}^2 \text{ s}^{-1}$</th></tr> </thead> <tbody> <tr> <td>150.0</td><td>20.0</td><td>60000</td></tr> <tr> <td>45.0</td><td>40.0</td><td>72000</td></tr> <tr> <td>22.5</td><td>60.0</td><td>81000</td></tr> <tr> <td>15.0</td><td>80.0</td><td>96000</td></tr> </tbody> </table>	R / s^{-1}	x / cm	$Rx^2 / \text{cm}^2 \text{ s}^{-1}$	150.0	20.0	60000	45.0	40.0	72000	22.5	60.0	81000	15.0	80.0	96000	3
R / s^{-1}	x / cm	$Rx^2 / \text{cm}^2 \text{ s}^{-1}$															
150.0	20.0	60000															
45.0	40.0	72000															
22.5	60.0	81000															
15.0	80.0	96000															
20(d)	<p>The tracks are thick indicating a heavily ionizing radiation (1) The tracks are straight indicating that the radiations are massive (1) Or the tracks are all about the same length so all the radiations have the same energy (1) Therefore the tracks are made by radiation from an alpha source [dependent on MP1 or MP2]</p>	3															
Total for question 20		12															

Question Number	Answer	Mark
21(a)	<p>Use of $\lambda_{\max} T = 2.898 \times 10^{-3} \text{ m K}$ (1)</p> <p>$T = 5800 \text{ (K)}$ (1)</p> <p><u>Example of calculation</u></p> $T = \frac{2.898 \times 10^{-3} \text{ m K}}{5.0 \times 10^{-7} \text{ m}} = 5796 \text{ K}$	2
21(b)	<p>Use of $I = \frac{L}{4\pi d^2}$ (1)</p> <p>Use of $L = \sigma AT^4$ (1)</p> <p>Use of $A = 4\pi r^2$ (1)</p> <p>$r = 7.0 \times 10^8 \text{ m}$ (ecf from (a)) (1)</p> <p><u>Example of calculation</u></p> $L = 590 \text{ W m}^{-2} \times 4\pi \times (2.3 \times 10^{11} \text{ m})^2 = 3.92 \times 10^{26} \text{ W}$ $A = \frac{3.92 \times 10^{26} \text{ W}}{5.67 \times 10^{-8} \text{ W m}^{-2} \text{ K}^{-4} \times (5800 \text{ K})^4} = 6.11 \times 10^{18} \text{ m}^2$ $r = \sqrt{\frac{6.11 \times 10^{18} \text{ m}^2}{4\pi}} = 6.97 \times 10^8 \text{ m}$	4
21(c)	<p>Use of 22% (1)</p> <p>Use of efficiency = $\frac{\text{useful energy output}}{\text{total energy input}}$ (1)</p> <p>Use of $I = P/A$ with $A = \pi r^2$ (1)</p> <p>$P = 1030 \text{ W}$ (1.03 kW) so the power requirement is met. (1)</p> <p><u>Example of calculation</u></p> $I = 0.78 \times 590 \text{ W m}^{-2} = 460 \text{ W m}^{-2}$ $A = \pi \times (1.1 \text{ m})^2 = 3.8 \text{ m}^2$ $P = 0.295 \times 460 \text{ W m}^{-2} \times 3.8 \text{ m}^2 \times 2 = 1030 \text{ W}$	4
Total for question 21		10

